

# Inhibitors of propagation of coagulation (factors VIII, IX and XI): a review of current therapeutic practice

Massimo Franchini<sup>1</sup> & Pier Mannuccio Mannucci<sup>2</sup>

<sup>1</sup>*Immunohaematology and Transfusion Centre, Department of Pathology and Laboratory Medicine, University Hospital of Parma, Parma and* <sup>2</sup>*Scientific Direction, IRCCS Cà Granda Foundation Maggiore Policlinico Hospital, Milan, Italy*

## Correspondence

Dr Pier Mannuccio Mannucci, Scientific Direction, IRCCS Cà Granda Foundation Maggiore Policlinico Hospital, Via Pace, 9, 20122 Milan, Italy.  
Tel.: + 39 02 5503 5422  
Fax: + 39 02 5032 0723  
E-mail: pmmannucci@libero.it

## Keywords

factor VIII, factor IX, factor XI, haemophilia, inhibitors, therapy

## Received

16 September 2010

## Accepted

13 December 2010

## Accepted Article

5 January 2011

The management of patients with congenital haemophilia who develop alloantibodies against factors of the propagation phase of blood coagulation, commonly known as inhibitors, is the most important challenge facing haemophilia caregivers at present, as this complication not only compromises the efficacy of replacement therapy but also consumes an enormous amount of economic resources. Development of inhibitors further complicates the clinical course of severe haemophilia, with a prevalence of up to 30% in patients with haemophilia A (factor VIII deficiency) and up to 5% in those with haemophilia B (factor IX deficiency) and haemophilia C (factor XI deficiency). While the short-term goal of treatment of patients who develop alloantibodies is the control of bleeding, the eradication of the inhibitor is the main long-term goal. The management of severe bleeding episodes and the eradication of the autoantibody are also the mainstays of treatment of patients with acquired haemophilia, a rare but life-threatening haemorrhagic condition characterized by the development of inhibitory autoantibodies against coagulation factor VIII. The most recent options available for treating patients with congenital haemophilia complicated by inhibitors and acquired haemophilia because of autoantibodies against factor VIII are summarized in this review article.

## Introduction

In patients with congenital haemophilia, the most serious and challenging complication of replacement therapy with coagulation factor concentrates is the development of alloantibodies, commonly known as inhibitors, that inhibit factors involved in the propagation phase of blood coagulation including factor VIII (FVIII, haemophilia A), factor IX (FIX, haemophilia B) or factor XI (FXI, haemophilia C) [1, 2]. These inhibitory alloantibodies develop in approximately 25–30% of severe haemophilia A patients and in 3–5% of patients with haemophilia B or C. The FVIII/IX Subcommittee of the International Society on Thrombosis and Haemostasis (ISTH) defines high-responder inhibitors as those with peak activity >5 Bethesda Units (BU) and with an anamnestic rise following replacement of the deficient clotting factor [3]. Inhibitory antibodies against coagulation factors may also rarely arise as autoantibodies in non-haemophilic persons [4]. In over 98% of the cases, the inhibitor is directed against coagulation FVIII, and this leads to the acquired form of haemophilia A [4]. Inhibitors render replacement therapy ineffective, thereby preclud-

ing patients from a safe and effective standard of care and predisposing them to a high risk of morbidity and mortality [5–8].

In this review, we summarize the most important laboratory and clinical characteristics of these allo- and autoantibodies, focusing mainly on their management.

## Inhibitors in congenital haemophilias

### General characteristics of inhibitors of factors VIII, IX and XI

Anti-FVIII alloantibodies are polyclonal, high-affinity immunoglobulins belonging to IgG subclasses, with IgG4 over-represented. They do not bind complement and usually react with the active sites of the FVIII molecule, primarily with epitopes in the protein domains A2 (resulting in the inhibition of FVIII-FIX complex formation) and C2 (inhibition of FVIII-von Willebrand factor and FVIII-phospholipid membrane interactions), and typically display type 1 reaction kinetics [9, 10]. A number of factors, both genetic (e.g. ethnicity, FVIII gene mutations, major

**Table 1**

Main characteristics of FVIII, FIX and FXI alloantibodies

Alloantibodies	Prevalence	Ig type	Protein domains*	Risk factors
<b>FVIII</b>	25–30%	IgG4	A2, C2	Genetic Genetic mutations (large deletions, nonsense mutations, <i>F8</i> gene inversions), inhibitor family history, ethnicity (African), immuno-regulatory genes (TNF- $\alpha$ , IL-10) Environmental Treatment-related factors (age at first factor concentrate exposure, intensive exposure to FVIII, source of FVIII product)
<b>FIX</b>	3–5%	IgG4	Gla, serine protease	Genetic mutations (large deletions, stop codons, frameshift mutations)
<b>FXI</b>	3–5%	IgG	Various functional domains of heavy and light chains	Glu117stop mutation (Type II mutation)

TNF, tumour necrosis factor; IL, interleukin; Ig, immunoglobulin; Gla,  $\gamma$ -carboxyglutamic acid region. \*Domains of the coagulation factor protein where epitopes of the inhibitor activity are located.

histocompatibility complex genotype, polymorphisms of immune-response genes) and environmental (e.g. number of FVIII exposure days, age at first exposure of FVIII concentrate, source of FVIII concentrate, modality and intensity of treatment) affect inhibitor formation, resulting in a complex multifactorial pathogenetic mechanism [11–16].

In haemophilia B, the prevalence of FIX inhibitors is 5–10 times lower than in haemophilia A [17]. FIX inhibitors are predominantly of the IgG4 subclass and have affinity for both the heavy and light chains of FIX. As in haemophilia A, some defects in the FIX gene (large deletions, stop codons and frameshift mutations) are associated with a greater likelihood of inhibitor development [18]. Unlike FVIII inhibitors, FIX inhibitors are accompanied in approximately half of the cases by severe anaphylactic reactions to the infusion of FIX-containing products [19, 20], which occur most often in children after relatively few exposure days (approximately 10–20) to any type of FIX-containing product [19]. FXI inhibitors are polyclonal IgG alloantibodies that act on various epitopes of the FXI molecule and inhibit activation of FXI by thrombin or FXIIa and activation of FIX by FXIa [6, 8]. Salomon *et al.* assessed the prevalence of inhibitors in 118 patients with severe FXI deficiency and found that 6% of them presented an inhibitor and that all were previously exposed to plasma replacement therapy [6]. Genotype analysis showed that all the patients with inhibitors were homozygous for the Glu117stop FXI gene mutation (so called type II mutation), which is accompanied by a baseline FXI < 1%. Thus, in patients carrying this type of mutation plasma-derived products containing FXI should be avoided as much as possible in order to prevent inhibitor formation. Table 1 summarizes the main characteristics of inhibitory alloantibodies to FVIII, FIX and FXI.

### Management of factor VIII, IX and XI inhibitors

While the immediate management of inhibitors consists of treating the acute bleeding event, long-term management has the goal to eradicate the inhibitor [21].

**Treatment of bleeding** In haemophiliacs with low-titre inhibitors (<5 BU), acute bleeding episodes can be controlled with high doses of FVIII, FIX or FXI concentrates, which can overcome the presence of inhibitors and allow the attainment of haemostatic levels of the factor infused [22]. The recommended bolus dosage corresponds to the sum of the inhibitor neutralizing dose plus the incremental dose (i.e. the usual therapeutic dose). The neutralizing dose is obtained by multiplying the inhibitor level by the plasma volume. If needed subsequent doses correspond to the incremental dose, administered either every 6–12 h as boluses or as a continuous intravenous infusion [23].

a) Use of bypassing agents Bypassing agents, such as activated prothrombin complex concentrates (APCCs) and recombinant activated factor VII (rFVIIa, NovoSeven, Novo Nordisk A/S, Bagsværd, Denmark) are indicated for patients with high-titre inhibitors (>5 BU) that do not respond to factor infusion [22–25]. The APCC Factor Eight Inhibitor Bypassing Agent or FEIBA (Baxter, Deerfield, IL, USA) is recommended at doses of 50–100 IU kg<sup>-1</sup> every 8–24 h, not exceeding 200 IU kg<sup>-1</sup> per day in order to decrease the risk of thrombotic events [26]. The optimal dosage of rFVIIa ranges from 90 to 120  $\mu$ g kg<sup>-1</sup> [27]. The cross-over study FENOC, FEIBA Novo Seven Comparative (FENOC), comparing these two bypassing agents in the treatment of acute bleeding episodes in haemophilia A patients with inhibitors showed a high success rate with both agents (80% for FEIBA and 78% for rFVIIa) but failed to reach the goal of equivalence [28]. The results of FENOC did show substantial within-individual discordance in the efficacy of both bypassing agents, as at the 2 h time point nearly half of the patients rated one product effective and the other ineffective in terms of haemostatic efficacy [28].

A recent systematic review of studies including haemophilia A and B patients with inhibitors concluded that the overall efficacy and bleeding control rates are higher for rFVIIa than for APCC (81–91% and 64–80%, respectively) when standard dosage regimens are used to treat mild-to-

moderate bleeds in inhibitor patients [29]. Another review, which used a Bayesian meta-regression model to evaluate the outcome of more than 2000 joint bleeds, found that the cumulative rate of control of bleeding at 12, 24 and 36 h was 66%, 88% and 95% for a standard rFVIIa regimen, but was lower for standard APCC therapy (39%, 62% and 76%). These differences were statistically significant and appeared robust in sensitivity analyses [30].

On the whole, there is substantial evidence that both bypassing agents are effective in controlling acute bleeding episodes, even though the success rate is sometimes lower than that of factor concentrate in patients without inhibitors. Both products have also a good safety profile with a low thrombotic risk [31] when used according to the approved indications in patients with bleeding disorders. On the other hand, the off-label use of rFVIIa is associated with a high risk of arterial thrombosis, especially among the elderly [32]. There is no evidence that either product is more efficacious than the other, but clinicians know that some patients may respond to one product and not to the other [2]. Because recombinant FVIIa does not contain FIX, this product is also the most suitable treatment choice for haemophilia B patients with inhibitors who developed anaphylactic reactions to infused FIX [3]. Finally, rFVIIa has also been successfully used for the management of bleeding unresponsive to antifibrinolytics in FXI deficient patients with inhibitors [33].

**b) High rFVIIa dosages** Recently, the use of rFVIIa in bolus doses larger than the standard doses mentioned above ( $90$  to  $120\text{ }\mu\text{g kg}^{-1}$ ) has been considered. Parameswaran *et al.* reported the results obtained in the frame of a retrospective registry of haemophilia A and B patients with inhibitors treated with various doses of rFVIIa, and reported an 84% response rate with doses  $<200\text{ }\mu\text{g kg}^{-1}$  and a 97% response rate with doses  $>200\text{ }\mu\text{g kg}^{-1}$  [34]. A prospective randomized trial compared a standard dose of rFVIIa ( $90\text{ }\mu\text{g kg}^{-1}$  repeated as necessary every 3 h) with a high single dose ( $270\text{ }\mu\text{g kg}^{-1}$ ) for home treatment of haemarthroses in 20 haemophiliacs with inhibitors [35]. The high-dosage rFVIIa regimen was effective, safe and required fewer rFVIIa infusions, thus simplifying home treatment. In a multicentre, randomized, double-blind, cross-over trial, Kavakli *et al.* [36] evaluated the efficacy and safety of two rFVIIa doses for treating haemarthroses in patients with congenital haemophilia A or B and inhibitors. Patients were randomly allocated to treat a first joint bleeding episode with one  $270\text{ }\mu\text{g kg}^{-1}$  rFVIIa dose, followed by two doses of placebo at 3 h intervals, and a second joint bleed with three single doses of  $90\text{ }\mu\text{g kg}^{-1}$  rFVIIa at 3 h intervals, or *vice versa*. Treatment was rated as effective for 65% of patients using the  $270\text{ }\mu\text{g kg}^{-1}$  dose vs. 70% for the  $90\text{ }\mu\text{g kg}^{-1} \times 3$  regimen. No safety issues were identified. Thus, the authors concluded that the administration of rFVIIa as a single  $270\text{ }\mu\text{g kg}^{-1}$  dose to treat haemarthroses in patients with haemophilia and inhibitors

was at least as efficacious and safe as the  $90\text{ }\mu\text{g kg}^{-1} \times 3$  regimen. Finally, another randomized study compared a single  $270\text{ }\mu\text{g kg}^{-1}$  bolus dose of rFVIIa vs. standard  $90\text{ }\mu\text{g kg}^{-1}$  doses of rFVIIa (a total of three doses were administered every 3 h) and  $75\text{ U kg}^{-1}$  dose of APCC (FEIBA) for the home management of joint bleeds. The authors observed a trend towards a better response with rFVIIa (successful responses were 37.5% with rFVIIa single high dose, 54.5% with rFVIIa standard dose and 27.3% with FEIBA standard dose), although differences were not statistically significant [37].

On the whole, these studies show that a single large dose of  $270\text{ }\mu\text{g kg}^{-1}$  rFVIIa is at least as effective as repeated smaller standard doses ( $90\text{ }\mu\text{g kg}^{-1}$ ), with obvious practical advantages in terms of conservation of the venous access particularly in children.

**Prophylaxis of bleeding** Bypassing agents are also increasingly being considered for secondary prophylaxis in patients with inhibitors, and early reports suggest a significant reduction of bleeding and an improvement of quality of life [38, 39]. At the moment there is only one published randomized trial of secondary prophylaxis with the rFVIIa administered with the goal to decrease the particularly high frequency of bleeding in 22 patients with haemophilia A [40]. After 3 months of secondary prophylaxis with two different daily doses of rFVIIa ( $90\text{ }\mu\text{g kg}^{-1}$  vs.  $270\text{ }\mu\text{g kg}^{-1}$ ), the patients enrolled in the study were followed for 3 additional months of on-demand treatment of bleeding episodes. There was a marked, statistically significant reduction of joint bleeding during prophylaxis, with a reduction, although non-statistically significant, of the frequency of haemarthrosis in patients randomized to the higher dose of  $270\text{ }\mu\text{g kg}^{-1}$  (59% reduction vs. 45% reduction for the  $90\text{ }\mu\text{g kg}^{-1}$  dosage). Surprisingly, the improvement somewhat persisted during the period of on-demand therapy, not only in terms of reduction of haemarthrosis frequency but also of more attendance at school or work [41]. A larger randomized study of secondary prophylaxis, carried out in unselected patients at an early time of inhibitor development, is currently starting (ENJOIH) [42].

On the whole, there is interest to use bypassing agents in inhibitor patients for regular prophylaxis, a method of treatment delivery that is so successful in children with uncomplicated haemophilia. Unfortunately the costs of secondary prophylaxis are huge [1], so there is a need to investigate further cost-effectiveness of this therapeutic strategy before its use can become widespread [43].

**Immunoadsorption** In the case of failure of bypassing agents to control bleeding, immunoadsorption may temporarily reduce the inhibitor titre in high-responder inhibitor patients, enabling effective replacement therapy with factor concentrates [44]. Freiburghaus *et al.* [45] reviewed the Malmö experience between 1980 and 1995 in 10 patients (five with haemophilia A and five with



haemophilia B) undergoing 19 procedures of immunoadsorption using staphylococcal protein A adsorption in a two-column system. In all but one case, inhibitor levels were dramatically lowered, allowing the subsequent maintenance of haemostatic levels of coagulation factors by means of factor concentrate therapy for a period long enough (usually 5 to 9 days) to stop ongoing bleeding or to prevent excessive bleeding at surgical interventions. Protein A immunoadsorption has also been successfully used in the context of immune tolerance induction (ITI) programmes (see below), in order to reduce inhibitor concentrations at the onset of treatment to levels low enough to allow delivery of a neutralizing dose of FVIII or FIX [46, 47].

Other positive experiences, during or not during ITI regimens, have been recorded using immunoadsorption of anti-FVIII alloantibodies to polyclonal sheep antibodies against human immunoglobulins [48].

*Inhibitor eradication by immune tolerance induction* This is the only strategy able to eradicate persistent inhibitors in severe haemophilia A patients [49]. Thirty years of ITI experience have shown global success rates ranging between 60% and 80% and helped to define the patients profile associated with higher success likelihood [50]. Most data on predictors of ITI success have been identified in the frame of three large retrospective registries: the International Immune Tolerance Registry (IITR), the North American Immune Tolerance Registry (NAITR) and the German Immune Tolerance Registry (GITR) [51–53]. The FVIII dose regimens include 200–300 IU kg<sup>-1</sup> day<sup>-1</sup> in the GITR and 50–200 IU kg<sup>-1</sup> day<sup>-1</sup> in the IITR and NAITR. Low inhibitor titre before ITI start (<10 BU) and lack of high anamnestic response (historical inhibitor peak titre <200 BU) were the most consistent predictors of ITI success. In the meta-analysis of data from the IITR and NAITR, low pre-ITI inhibitor titre (<10 BU) was also associated with a more rapid tolerance [54]. However, the limits of these registries were that they retrospectively collected data differing from each other in terms of patient and treatment characteristics and particularly on the definition of end points.

In the last few years, two large-scale randomized trials, the International Immune Tolerance Induction study (I-TI) [55] and the Rescue Immunotolerance study (RESIST) [56], have been conducted with the aim to resolve some of the unanswered questions on ITI, that is chiefly the optimal ITI regimen. However, the I-TI study, aimed at evaluating the success rate and time to success in 150 patients with good prognostic profile (historical peak titre ≥5 BU and ≤200 BU; starting titre <10 BU) randomized to receive FVIII doses of 50 IU kg<sup>-1</sup> three times weekly or 200 IU kg<sup>-1</sup> daily, was interrupted prematurely because of safety concerns. A significantly greater cumulative number of bleeding episodes in joint and non-joint sites was indeed observed in the low-dose arm vs. the high-dose arm, at all stages of ITI, but particularly in the first ITI phase, when inhibitors were still detectable [57]. At study termination, although no

sufficient power proving therapeutic equivalence was reached, ITI success rates were not different in the two treatment arms. However, median time to achieve negative inhibitor titre and normal FVIII recovery were significantly shorter (about 50%) in patients receiving the high-dose regimen. Thus, the earlier attainment of tolerance with the higher dosage (and hence of measurable FVIII levels in plasma) perhaps explains the difference in the number of bleeding episodes between the two arms.

There are fewer published reports of ITI in haemophilia B, given the fact that FIX inhibitors are rarer. At variance with results in patients with FVIII inhibitors, the success of ITI in eradicating FIX inhibitors is low, particularly in patients who develop anaphylactic reactions [25]. Among 16 haemophilia B patients in the NAITR who completed ITI, five had a successful outcome using FIX dosing regimens that ranged from 43 to 200 IU kg<sup>-1</sup> day<sup>-1</sup> [53]. Adverse events were reported during 11 of 17 (65%) ITI courses, a frequency 10 times higher than that for persons with FVIII inhibitors. Anaphylactic reactions accounted for 11 of the 14 adverse events and represented the major reason for failure in at least four of 11 unsuccessful courses of ITI in haemophilia B. Moreover, an association between those reactions to FIX in patients undergoing ITI and the development of the nephrotic syndrome was observed, because three of the 10 patients with those reactions developed this renal complication. The ISTH Registry on FIX inhibitors data showed that ITI was successful in only five of the 34 patients (15%) in whom it was attempted, and two of these patients had low-responder inhibitors [8]. Moreover, 13 of the 34 patients (38%) developed the nephrotic syndrome during ITI.

With the aim to provide useful information to haemophilia caregivers, an international panel of experts developed consensus recommendations for ITI, with ratings based on the level of supporting evidence [49]. For haemophilia A, the initiation of ITI was recommended after the inhibitor titre dropped to <10 BU (level IIb). While among good-risk patients (i.e. historical peak titre <200 BU, pre-ITI titre <10 BU, <5 years since diagnosis) no dosing regimen is clearly superior to another, among poor-risk patients (i.e. historical peak titre >200 BU, pre-ITI titre >10 BU, >5 years since diagnosis) a higher success rate was obtained with the use of high-dose regimens (≥200 IU kg<sup>-1</sup> day<sup>-1</sup>; level IIb). The consensus recommendations on the FVIII product type were that ITI is successful using FVIII products regardless of VWF content, with no definitive evidence supporting the superiority of any FVIII product, and that most patients can be effectively tolerized with the same FVIII product in use at the time of inhibitor detection (level IIb).

For haemophilia B, the panel concluded that evidence was insufficient to recommend ITI or to establish predictors for success or failure to ITI. If a decision to start ITI is made, the inclusion of a routine urinalysis is recommended in the follow-up in order to detect proteinuria and early nephrotic syndrome (level IV). As for FIX inhibitors, there are no solid data on the responsiveness to ITI in patients

**Table 2**

Therapeutic options for haemophilia patients with FVIII, FIX and FXI alloantibodies

<b>Treatment of bleeding</b>	
<b>Low-titre inhibitors (&lt;5 BU)</b>	High doses of factor concentrates
<b>High-titre inhibitors (&gt;5 BU)</b>	
<b>First line</b>	rFVIIa, APCC
<b>Second line</b>	Immunoadsorption
<b>Inhibitor eradication</b>	
<b>First line</b>	Immune tolerance induction*
<b>Second line</b>	Rituximab

rFVIIa, recombinant activated factor VII; APCC, activated prothrombin complex concentrates. \*The immune tolerance induction success in eradicating FIX and FXI alloantibodies is low.

with a persistent FXI inhibitor [58]. Table 2 summarizes the therapeutic options for the management of FVIII, FIX and FXI alloantibodies.

Finally, it is important to mention that some recent data suggest that the addition of rituximab (a monoclonal antibody against CD20-positive B cells) may improve response rates in patients with incomplete or no response to the ITI regimen. However, published experience with this agent in the treatment of alloantibodies associated with congenital haemophilia is limited to case reports or small case series [59].

## Acquired haemophilia

### General characteristics of acquired factor VIII inhibitors

Acquired inhibitors against FVIII occur rarely in the non-haemophilic population, with an incidence of between 1.3 and 1.5 cases per million annually [60–62]. Although uncommon, these autoantibodies are associated with a high rate of morbidity and mortality, because severe bleeds occur in up to 90% of affected patients and the mortality rate is high, ranging from 8% to 22% [63, 64]. The age distribution of autoantibodies is typically biphasic with a small peak between 20 and 30 years, because of post-partum inhibitors, and a major peak in elderly patients. The incidence in men and women is similar, except in the age ranges 20–40 years when the effect of pregnancy leads to a higher prevalence in women. In approximately 50% of cases, factor VIII autoantibodies occur in patients with no concomitant disease, while the remaining cases may be associated with the post-partum period, autoimmune diseases (systemic lupus erythematosus, rheumatoid arthritis, thyroid disorders), haematological or solid cancers, infections or use of medications [65–68].

The bleeding pattern of acquired haemophilia is rather different from that of congenital haemophilia A. Most patients with FVIII autoantibodies have haemorrhages into

**Table 3**

Therapeutic options for acquired haemophilia A

<b>Antihemorrhagic treatment</b>	
<b>First line</b>	rFVIIa, APCC
<b>Alternative treatment*</b>	FVIII concentrates, DDAVP
<b>Second line</b>	Immunoadsorption or plasmapheresis
<b>Inhibitor eradication</b>	
<b>First line</b>	Corticosteroids ± cyclophosphamide
<b>Second line</b>	Rituximab ± corticosteroids
<b>Alternative treatment</b>	Azathioprine, vincristine, mycophenolate, cyclosporin, intravenous immunoglobulin

rFVIIa, recombinant activated factor VII; APCC, activated prothrombin complex concentrates; DDAVP, desmopressin. \*Low-titre inhibitors and minor bleeding.

the skin, muscles or soft tissues and mucous membranes (e.g. epistaxis, gastrointestinal and urological bleeds, retro-peritoneal haematomas, post-partum bleeding), whereas haemarthroses, a typical feature of congenital FVIII deficiency, are uncommon [62]. The haemorrhages are often serious and the disease may manifest more dramatically by life-threatening bleeding following trauma or surgery or by cerebral haemorrhage.

Factor VIII autoantibodies are mostly IgG1 and IgG4 autoantibodies, often acting with second-order kinetics and reacting with the same regions of the FVIII molecule targeted by alloantibodies (i.e. A2 and C2 domains). The diagnosis of acquired haemophilia in a patient with no previous personal or family history of bleeding is typically based upon the initial detection of an isolated prolongation of the activated partial thromboplastin time (APTT), which cannot be corrected by incubating for 2 h at 37°C equal volumes of patient and normal plasma, and subsequent identification of a low FVIII level with evidence of FVIII inhibitory activity (titrated using the Bethesda method) [4, 69].

### Management of acquired factor VIII inhibitors

Similar to the treatment of FVIII alloantibodies, the management of acquired haemophilia A is directed to the control of bleeding episodes and the eradication of the inhibitor [70] (Table 3).

*Treatment of bleeding* Efficient haemostasis can be achieved with a variety of methods that, if necessary, may be used in combination: normalization/correction of FVIII deficiency (human plasma derived or recombinant FVIII concentrates, desmopressin), bypassing the inhibitor activity (FEIBA or rFVIIa), neutralization of the inhibitor by idiotypic anti-FVIII antibodies (high-dose immunoglobulin) and removal of the inhibitor by plasmapheresis or immunoadsorption. The choice of the most appropriate therapeutic strategy depends on the site and severity of the haemorrhage, patient characteristics, underlying disorder and

inhibitor titre [71,72]. FVIII replacement therapy is the treatment of choice when bleeding is minor and the inhibitor titre low (<5 BU). A loading dose is given as bolus to neutralize the inhibitor and to achieve haemostatic FVIII levels, followed by maintenance doses given by bolus or continuous infusion. The experience with desmopressin in acquired haemophilia is very limited and available data indicate that it may be useful only in patients with a low-titre inhibitor for the treatment of minor bleeding episodes [73].

Bypassing agents are recommended as first-line treatment for severe bleeds and both rFVIIa and FEIBA are effective, although there are no comparative trials to demonstrate superior efficacy for either product [74]. Sallah retrospectively analysed the efficacy of FEIBA in 34 acquired haemophilia patients and reported 100% haemostatic efficacy for moderate bleeds and 75% for severe bleeds, with an overall response rate of 86% [75]. In a retrospective analysis of 38 patients, Hay *et al.* [76] reported a positive response in 100% of patients when rFVIIa was used as a first-line treatment, and a positive response in 75% of patients when it was used as salvage therapy after failure of therapy with blood products.

Recently, Sumner *et al.* [77] collected the available data on the compassionate use of rFVIIa in acquired haemophilia patients from the Haemophilia and Thrombosis Research Society (HTRS) registry and the published literature. A total of 139 patients were treated with rFVIIa for 204 bleeding episodes. The overall success rate (complete or partial) of rFVIIa was 88% (161/182 bleeding episodes evaluable). rFVIIa as a first-line treatment was effective overall in 95% of bleeding episodes compared with 80% when it was used as salvage therapy after failure of other haemostatic agents.

High-dose immunoglobulins have also been used in acquired haemophilia. In a prospective multicentre study, 19 patients with low-titre inhibitors were treated with 400 mg kg<sup>-1</sup> for 5 consecutive days or 1000 mg kg<sup>-1</sup> for 2 days, with an overall response rate of 25% [78]. Thus, high-dose immunoglobulins are not the first choice for the eradication of FVIII autoantibodies, but may play a role as adjunctive therapy to other inhibitor eradicating treatments (steroids, immunoadsorption, immune tolerance regimens) [70]. In patients with a high-titre inhibitor and severe haemorrhages, the extracorporeal removal of the autoantibody by therapeutic plasmapheresis or its immunoadsorption to staphylococcal protein A or to polyclonal sheep antibodies against human immunoglobulins can be used prior to factor concentrate treatment [44].

**Inhibitor eradication** In acquired haemophilia, inhibitor eradication may be obtained with immunosuppressive agents including corticosteroids and drugs such as cyclophosphamide, azathioprine, 6-mercaptopurine and vincristine [79–83]. In their meta-analysis combining data from 20 reports, Delgado *et al.* [61] concluded that cyclophosphamide was superior to prednisone in terms of inhibitor eradi-

cation, but not in terms of overall survival. The combined data from uncontrolled cohort studies recently reviewed by Collins [72] suggested a benefit for combined steroids and cytotoxic agents. More recently, biotherapy with rituximab has also been used to treat patients with acquired haemophilia, with high success rates [84, 85]. A literature review collecting 65 patients with acquired haemophilia A treated with rituximab found that a complete or partial response was reached in more than 90% of cases [86]. Finally, preliminary data from ongoing studies have supported the effectiveness of ITI protocols in eradicating inhibitors also in patients with acquired haemophilia [87, 88].

## Conclusions

In the last two decades, remarkable progress has been made in the management of patients with inhibitors of coagulation factors of the propagation phase of blood coagulation. In particular, the introduction of bypassing agents has dramatically improved the management of acute bleeding, allowing home treatment with a substantial amelioration of patient quality of life. At the same time, the widespread implementation of ITI has permitted inhibitor eradication in an increasing number of patients, so that the life expectancy in inhibitor patients has become similar to that of severe haemophiliacs without inhibitors [89, 90]. Pertaining to innovative therapies, rituximab has been shown to be a promising agent for ITI-resistant patients, which warrants further investigation in large prospective studies. This agent is indeed a strong candidate therapy for the eradication of autoantibodies in acquired haemophilia.

Finally, efforts are currently being made to reveal more about the pathophysiology of the development of inhibitors and of possible therapeutic strategies. For instance, an ongoing randomized trial (SIPPET study) is tackling the issue of whether or not there is a difference in inhibitor formation in congenital haemophilia between children previously untreated or minimally treated with plasma-derived, VWF-containing FVIII products, compared with children treated exclusively with recombinant FVIII [91, 92]. Ultimately, future insights into the understanding of the immunobiology of inhibitor formation will unlock the key to the design of more tailored and cost-effective therapies for the management and prevention of inhibitors.

## Competing Interests

MF has no conflict of interest. PMM has received speaker honoraria from Baxter and Novo Nordisk, the manufacturers of FEIBA and Novo Seven.

*There was no financial support for the content of this review article.*

## REFERENCES

- 1 Gringeri A, Mantovani LG, Scalone L, Mannucci PM. Cost of care and quality of life for patients with hemophilia complicated by inhibitors: the COCIS Study Group. *Blood* 2003; 102: 2358–63.
- 2 Haya S, Moret A, Cid AR, Cortina V, Casaña P, Cabrera N, Aznar JA. Inhibitors in haemophilia A: current management and open issues. *Haemophilia* 2007; 13 (Suppl. 5): 52–60.
- 3 White GC, Rosendaal F, Aledort LM, Lusher JM, Rothschild C, Ingerslev J, Factor VIII, Factor IX Subcommittee. Definitions in hemophilia. Recommendations of the scientific subcommittee on factor VIII and factor IX of the scientific and standardization committee of the International Society on Thrombosis and Haemostasis. *Thromb Haemost* 2001; 85: 560.
- 4 Franchini M, Targher G, Montagnana M, Lippi G. Laboratory, clinical and therapeutic aspects of acquired hemophilia A. *Clin Chim Acta* 2008; 395: 14–8.
- 5 Lusher JM. Inhibitor antibodies to factor VIII and factor IX: management. *Semin Thromb Hemost* 2000; 26: 179–88.
- 6 Salomon O, Zivelin A, Livnat T, Dardik R, Loewenthal R, Avishai O, Steinberg DM, Rosove MH, O'Connell N, Lee CA, Seligsohn U. Prevalence, causes, and characterization of factor XI inhibitors in patients with inherited factor XI deficiency. *Blood* 2003; 101: 4783–8.
- 7 Wight J, Paisley S. The epidemiology of inhibitors in haemophilia A: a systematic review. *Haemophilia* 2003; 9: 418–35.
- 8 Key NS. Inhibitors in congenital coagulation disorders. *Br J Haematol* 2004; 127: 379–91.
- 9 Scandella DH. Properties of anti-factor VIII antibodies in hemophilia A patients. *Semin Thromb Hemost* 2000; 26: 137–42.
- 10 Spiegel PC, Jacquemin M, Saint-Remy JM, Stoddard BL, Pratt KP. Structure of a factor VIII C2 domain-immunoglobulin G4k Fab complex: identification of an inhibitory antibody epitome on the surface of factor VIII. *Blood* 2001; 98: 13–9.
- 11 Cox-Gill J. The role of genetics in inhibitor formation. *Thromb Haemost* 1999; 82: 500–4.
- 12 Santagostino E, Mancuso ME, Rocino A, Mancuso G, Mazzucconi MG, Tagliaferri A, Messina M, Mannucci PM. Environmental risk factors for inhibitor development in children with haemophilia A: a case-control study. *Br J Haematol* 2005; 130: 422–7.
- 13 Lee CA, Lillicrap D, Astermark J. Inhibitor development in hemophiliacs: the roles of genetic versus environmental factors. *Semin Thromb Hemost* 2006; 32 (Suppl. 2): 10–4.
- 14 Gouw SC, van der Bom JG, Auerswald G, Ettinghausen CE, Tedgård U, van den Berg HM. Recombinant versus plasma-derived factor VIII products and the development of inhibitors in previously untreated patients with severe hemophilia A: the CANAL cohort study. *Blood* 2007; 109: 4693–7.
- 15 Gouw SC, van den Berg HM. The multifactorial etiology of inhibitor development in hemophilia: genetics and Environment. *Semin Thromb Hemost* 2009; 35: 723–34.
- 16 Iorio A, Halimeh S, Holzhauser S, Goldenberg N, Marchesini E, Marcucci M, Young G, Bidlingmaier C, Brandao LR, Ettingshausen CE, Gringeri A, Kenet G, Knöfler R, Kreuz W, Kurnik K, Manner D, Santagostino E, Mannucci PM, Nowak-Göttl U. Rate of inhibitor development in previously-untreated hemophilia A patients treated with plasma-derived or recombinant factor VIII concentrates. A systematic review. *J Thromb Haemost* 2010; 8: 1256–65.
- 17 Katz J. Prevalence of FIX inhibitors among patients with hemophilia B: results of large scale North American survey. *Haemophilia* 1996; 2: 38–41.
- 18 Ljung RC. Gene mutations and inhibitor formation in patients with haemophilia B. *Acta Haematol* 1995; 94 (Suppl. 1): 49–52.
- 19 Warrier I. Management of haemophilia B patients with inhibitors and anaphylaxis. *Haemophilia* 1998; 4: 574–6.
- 20 Jadhav M, Warrier I. Anaphylaxis in patients with hemophilia. *Semin Thromb Hemost* 2000; 26: 205–8.
- 21 Von Depka M. Managing acute bleeds in the patient with haemophilia and inhibitors: options, efficacy and safety. *Haemophilia* 2005; 11 (Suppl. 1): 18–23.
- 22 Gringeri A, Mannucci PM, for the Italian Association of Haemophilia Centers. Italian guidelines for the diagnosis and treatment of patients with haemophilia and inhibitors. *Haemophilia* 2005; 11: 611–9.
- 23 Schulman S. Continuous infusion. *Haemophilia* 2003; 9: 368–75.
- 24 Astermark J. Treatment of the bleeding inhibitor patient. *Semin Thromb Hemost* 2003; 29: 77–85.
- 25 Hay CR, Brown S, Collins PW, Keeling DM, Liesner R. The diagnosis and management of factor VIII and IX inhibitors: a guideline from the United Kingdom Haemophilia Centre Doctors Organisation. *Br J Haematol* 2006; 133: 591–605.
- 26 Ehrlich HJ, Enzl MJ, Gomperts ED. Safety of factor VIII inhibitor bypass activity (FEIBA): 10-year compilation of thrombotic adverse events. *Haemophilia* 2002; 8: 83–90.
- 27 Franchini M, Lippi G. Recombinant activated factor VII: mechanisms of action and current indications. *Semin Thromb Hemost* 2010; 36: 485–92.
- 28 Astermark J, Donfield SM, DiMichele DM, Gringeri A, Gilbert SA, Waters J, Berntorp E, FENOC Study Group. A randomized comparison of bypassing agents in hemophilia complicated by an inhibitor: the FEIBA NovoSeven Comparative (FENOC) Study. *Blood* 2007; 109: 546–51.
- 29 Knight C, Danø AM, Kennedy-Martin T. Systematic review of efficacy of rFVIIa and aPCC treatment for hemophilia patients with inhibitors. *Adv Ther* 2009; 26: 68–88.
- 30 Treur MJ, McCracken F, Heeg B, Joshi AV, Botteman MF, De Charro F, van Hout B. Efficacy of recombinant activated



- factor VII vs. activated prothrombin complex concentrate for patients suffering from haemophilia complicated with inhibitors: a Bayesian meta-regression. *Haemophilia* 2009; 15: 420–36.
- 31** Iorio A, Matino D, D'Amico R, Makris M. Recombinant factor VIIa concentrate versus plasma derived concentrates for the treatment of acute bleeding episodes in people with haemophilia and inhibitors. *Cochrane Database Syst Rev* 2010; 8: CD004449.
- 32** Levi M, Levy JH, Andersen HF, Truloff D. Safety of recombinant activated factor VII in randomized clinical trials. *N Engl J Med* 2010; 363: 1791–800.
- 33** Lawler P, White B, Pye S, Hermans C, Riddel A, Costello C, Brown S, Lee CA. Successful use of recombinant factor VIIa in a patient with inhibitor secondary to severe factor XI deficiency. *Haemophilia* 2002; 8: 145–8.
- 34** Parameswaran R, Shapiro AD, Gill JC, Kessler CM, HTRS Registry Investigators. Dose effect and efficacy of rFVIIa in the treatment of haemophilia patients with inhibitors: analysis from the Hemophilia and Thrombosis Research Society Registry. *Haemophilia* 2005; 11: 100–6.
- 35** Santagostino E, Mancuso ME, Rocino A, Mancuso G, Scaraggi F, Mannucci PM. A prospective randomized trial of high and standard dosages of recombinant factor VIIa for treatment of hemarthroses in hemophiliacs with inhibitors. *J Thromb Haemost* 2006; 4: 367–71.
- 36** Kavakli K, Makris M, Zulfikar B, Erhardtsen E, Abrams ZS, Kenet G, NovoSeven trial (F7HAEM-1510) investigators. Home treatment of hemarthroses using a single dose regimen of recombinant activated factor VII in patients with haemophilia and inhibitors. A multi-centre, randomised, double-blind, cross-over trial. *Thromb Haemost* 2006; 95: 600–5.
- 37** Young G, Shafer FE, Rojas P, Seremetis S. Single 270 microg kg-dose rFVIIa vs. standard 90 microg kg-dose rFVIIa and APCC for home treatment of joint bleeds in haemophilia patients with inhibitors: a randomized comparison. *Haemophilia* 2008; 14: 287–94.
- 38** Fischer K, Valentino L, Ljung R, Blanchette V. Prophylaxis for severe hemophilia: clinical challenges in the absence as well as in the presence of inhibitors. *Haemophilia* 2008; 14 (Suppl. 3): 196–201.
- 39** Franchini M, Manzato F, Salvagno GL, Montagnana M, Zaffanello M, Lippi G. Prophylaxis in congenital hemophilia with inhibitors: the role of recombinant activated factor VII. *Semin Thromb Hemost* 2009; 35: 814–9.
- 40** Konkle BA, Ebbesen LS, Erhardtsen E, Bianco RP, Lissitchkov T, Rusen L, Serban MA. Randomized, prospective clinical trial of recombinant factor VIIa for secondary prophylaxis in hemophilia patients with inhibitors. *J Thromb Haemost* 2007; 5: 1904–13.
- 41** Hoots WK, Ebbesen LS, Konkle BA, Auerswald GK, Roberts HR, Weatherall J, Ferran JM, Ljung RC, Novoseven (F7HAEM-1505) Investigators. Secondary prophylaxis with recombinant activated factor VII improves health-related quality of life of haemophilia patients with inhibitors. *Haemophilia* 2008; 14: 466–75.
- 42** Santagostino E, Awerswald G, Jimenez-Yuste V, Ljung R, Morfini M, Rocino A. European initiative to prevent joint damage in haemophilia children with inhibitor (ENJOIH)-a prospective study. *Haemophilia* 2010; 16 (Suppl. 4): 78. (abstract 17P63).
- 43** Mannucci PM, Palhares de Miranda PA. International survey of attitudes towards secondary prophylaxis with recombinant factor VIIa in haemophilia A patients with inhibitors. *Haemophilia* 2009; 15: 345–7.
- 44** Franchini M, Sassi M, Dell'Anna P, Manzato F, Salvagno GL, Montagnana M, Zaffanello M, Targher G, Lippi G. Extracorporeal immunoadsorption for the treatment of coagulation inhibitors. *Semin Thromb Hemost* 2009; 35: 76–80.
- 45** Freiburghaus C, Berntorp E, Ekman M, Gunnarsson M, Kjellberg BM, Nilsson IM. Immunoadsorption for removal of inhibitors: update on treatments in Malmö-Lund between 1980 and 1995. *Haemophilia* 1998; 4: 16–20.
- 46** Nilsson IM, Berntorp E, Zettervall O. Induction of immune tolerance in patients with hemophilia and antibodies to factor VIII by combined treatment with intravenous IgG, cyclophosphamide, and factor VIII. *N Engl J Med* 1988; 318: 947–50.
- 47** Freiburghaus C, Berntorp E, Ekman M, Gunnarsson M, Kjellberg B, Nilsson IM. Tolerance induction using the Malmö treatment model 1982–1995. *Haemophilia* 1999; 5: 32–9.
- 48** Jansen M, Schmaldienst S, Banyai S, Quehenberger P, Pabinger I, Defler K, Hörl WH, Knöbl P. Treatment of coagulation inhibitors with extracorporeal immunoadsorption (Ig-Therasorb). *Br J Haematol* 2001; 112: 91–7.
- 49** Di Michele D, Hoots WK, Pipe SW, Rivard GE, Santagostino E. International workshop on immune tolerance induction: consensus recommendations. *Haemophilia* 2007; 13 (Suppl. 1): 1–22.
- 50** Wight J, Paisley S, Knight C. Immune tolerance induction in patients with haemophilia A with inhibitors. *Haemophilia* 2003; 9: 436–63.
- 51** Lenk H. The German Registry of immune tolerance treatment in hemophilia – 1999 update. *Haematologica* 2000; 10 (Suppl.): 45–7.
- 52** Mariani G, Kroner B, for the Immune Tolerance Study Group. Immune tolerance in hemophilia with inhibitors: predictors of success. *Haematologica* 2001; 86: 1186–93.
- 53** Di Michele D. The North American Immune Tolerance Registry: contributions to the thirty-year experience with immune tolerance therapy. *Haemophilia* 2009; 15: 320–8.
- 54** Kroner BL. Comparison of the international immune tolerance registry and the North American immune tolerance registry. *Vox Sang* 1999; 77 (Suppl. 1): 33–7.
- 55** Di Michele DM, Hay CR. The international immune tolerance study: a multicenter prospective randomized trial in progress. *J Thromb Haemost* 2006; 4: 2271–3.
- 56** Gringeri A. VWF/FVIII concentrates in high-risk immunotolerance: the RESIST study. *Haemophilia* 2007; 13 (Suppl. 5): 73–7.



- 57 DiMichele DM, Goldberg I, Foulkes M, Hay CRM, on behalf of the International Immune Tolerance Study Group. International prospective randomized immune tolerance (ITI) study: preliminary results of therapeutic efficacy and safety [abstract 07S03]. *Haemophilia* 2010; 16 (Suppl. 4): 29.
- 58 Salomon O, Zivelin A, Livnat T, Seligsohn U. Inhibitors to factor XI in patients with severe factor XI deficiency. *Semin Hematol* 2006; 43 (Suppl. 1): S10–2.
- 59 Franchini M, Mengoli C, Lippi G, Targher G, Montagnana M, Salvagno GL, Zaffanello M, Cruciali M. Immune tolerance with rituximab in congenital haemophilia with inhibitors: a systematic literature review based on individual patients' analysis. *Haemophilia* 2008; 14: 903–12.
- 60 Cohen AJ, Kessler CM. Acquired inhibitors. *Baillieres Clin Haematol* 1996; 9: 331–54.
- 61 Delgado J, Jimenez-Yuste V, Hernandez-Navarro F, Villar A. Acquired haemophilia: review and meta-analysis focused on therapy and prognostic factors. *Br J Haematol* 2003; 121: 21–35.
- 62 Franchini M, Gandini G, Di Paolantonio T, Mariani G. Acquired hemophilia A: a concise review. *Am J Hematol* 2005; 80: 55–63.
- 63 Collins P, Macartney N, Davies B, Lees S, Giddings J, Maier R. A population based, unselected, consecutive cohort of patients with acquired haemophilia A. *Br J Haematol* 2004; 124: 86–90.
- 64 Collins PW, Percy CL. Advances in the understanding of acquired haemophilia A: implications for clinical practice. *Br J Haematol* 2010; 148: 183–94.
- 65 Green D, Lechner K. A survey of 215 non-hemophilic patients with inhibitors to factor VIII. *Thromb Haemost* 1981; 45: 200–3.
- 66 Yee TT, Pasi KJ, Lilley PA, Lee CA. Factor VIII inhibitors in haemophiliacs: a single-centre experience over 34 years, 1964–97. *Br J Haematol* 1999; 104: 909–14.
- 67 Sallah S, Wan JY. Inhibitors against factor VIII in patients with cancer. *Cancer* 2001; 91: 1067–74.
- 68 Franchini M. Postpartum acquired factor VIII inhibitors. *Am J Hematol* 2006; 81: 768–73.
- 69 Kershaw G, Jayakodi D, Dunkley S. Laboratory identification of factor inhibitors: the perspective of a large tertiary hemophilia centre. *Semin Thromb Hemost* 2009; 35: 760–8.
- 70 Franchini M, Lippi G. Acquired factor VIII inhibitors. *Blood* 2008; 112: 250–5.
- 71 Kessler CM. Acquired factor VIII autoantibody inhibitors: current concepts and potential therapeutic strategies for the future. *Haematologica* 2000; 85 (Suppl. 1): 57–63.
- 72 Collins PW. Treatment of acquired hemophilia A. *J Thromb Haemost* 2007; 5: 893–900.
- 73 Mudad R, Kane WH. DDAVP in acquired haemophilia A: case report and review of the literature. *Am J Haematol* 1993; 43: 295–9.
- 74 Huth-Kühne A, Baudo F, Collins P, Ingerslev J, Kessler CM, Lévesque H, Castellano ME, Shima M, St-Louis J. International recommendations on the diagnosis and treatment of patients with acquired hemophilia A. *Haematologica* 2009; 94: 566–75.
- 75 Sallah S. Treatment of acquired haemophilia with factor eight inhibitor bypassing activity. *Haemophilia* 2004; 10: 169–73.
- 76 Hay CRM, Negrier C, Ludlam CA. The treatment of bleeding in acquired hemophilia with recombinant factor VIIa: a multicenter study. *Thromb Haemost* 1997; 78: 3–7.
- 77 Sumner MJ, Geldziler BD, Pedersen M, Seremetis S. Treatment of acquired haemophilia with recombinant activated FVII: a critical appraisal. *Haemophilia* 2007; 13: 451–61.
- 78 Schwartz RS, Gabriel DA, Aledort LM, Green D, Kessler CM. A prospective study of treatment of acquired (autoimmune) factor VIII inhibitors with high-dose intravenous gammaglobulin. *Blood* 1995; 86: 797–804.
- 79 Green D. Suppression of an antibody to factor VIII by a combination of factor VIII and cyclophosphamide. *Blood* 1971; 37: 381–7.
- 80 Green D, Rademaker AW, Briet E. A prospective, randomized trial of prednisone and cyclophosphamide in the treatment of patients with factor VIII autoantibodies. *Thromb Haemost* 1993; 70: 753–7.
- 81 Petrovic M, Derom E, Baele G. Cyclosporine treatment of acquired hemophilia due to factor VIII antibodies. *Haematologica* 2000; 85: 895–6.
- 82 Shaffer LG, Phillips MD. Successful treatment of acquired hemophilia with oral immunosuppressive therapy. *Ann Intern Med* 1997; 127: 206–9.
- 83 Spero JA, Lewis JH, Hasiba U. Corticosteroid therapy for acquired F VIII:C inhibitors. *Br J Haematol* 1981; 48: 635–42.
- 84 Stasi R, Brunetti M, Stipa E, Amadori S. Selective B-cell depletion with rituximab for the treatment of patients with acquired hemophilia. *Blood* 2004; 103: 4424–8.
- 85 Wiestner A, Cho HJ, Asch AS, Michelis MA, Zeller JA, Peerschke EI, Weksler BB, Schechter GP. Rituximab in the treatment of acquired factor VIII inhibitor. *Blood* 2002; 100: 3426–8.
- 86 Franchini M. Rituximab in the treatment of adult acquired hemophilia A: a systematic review. *Crit Rev Oncol Hematol* 2007; 63: 47–52.
- 87 Nemes L, Pitlik E. Ten years experience with immunotolerance induction therapy in acquired hemophilia. *Haematologica* 2003; 88 (Suppl. 12): 106–10.
- 88 Zeitler H, Ulrich-Merzenich G, Hess L, Konsek E, Unkrig C, Walger P, Vetter H, Brackmann HH. Treatment of acquired hemophilia by the Bonn–Malmö Protocol: documentation of an in vivo immunomodulating concept. *Blood* 2005; 105: 2287–93.
- 89 Mannucci PM. Back to the future: a recent history of hemophilia treatment. *Haemophilia* 2008; 14 (Suppl. 3): 10–8.

- 90** Astermark J, Santagostino E, Keith Hoots W. Clinical issues in inhibitors. *Haemophilia* 2010; 16 (Suppl. 5): 54–60.
- 91** Study NCT 01064284; EUDRACT n. 2009-011186-88SIPPET study (SIPPET study). Available at <http://www.clinicaltrials.gov> (last accessed 10 December 2010).
- 92** Mannucci PM, Gringeri A, Peyvandi F, Santagostino E. Factor VIII products and inhibitor development: the SIPPET study (survey of inhibitors in plasma-product exposed toddlers). *Haemophilia* 2007; 13 (Suppl. 5): 65–8.